

Probing New Physics via an Angular Analysis of $B \rightarrow V_1 V_2$ Decays

David London ^{a,1}, Nita Sinha ^{b,2} and Rahul Sinha ^{b,3}

*a: Laboratoire René J.-A. Lévesque, Université de Montréal,
C.P. 6128, succ. centre-ville, Montréal, QC, Canada H3C 3J7*

*b: Institute of Mathematical Sciences, C. I. T Campus,
Taramani, Chennai 600 113, India*

(February 1, 2008)

Abstract

We show that an angular analysis of $B \rightarrow V_1 V_2$ decays yields numerous tests for new physics (NP) in the decay amplitudes. Many of these NP observables are nonzero even if the strong phase differences vanish. For certain observables, neither time-dependent measurements nor tagging is necessary. Should a signal for new physics be found, one can place a lower limit on the size of the NP parameters, as well as on their effect on the measurement of the phase of $B^0-\bar{B}^0$ mixing.

¹london@lps.umontreal.ca

²nita@imsc.res.in

³sinha@imsc.res.in

CP violation in the standard model (SM) is due to the presence of a nonzero complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix. This explanation can be tested by measuring CP-violating rate asymmetries in B decays, and extracting α , β and γ , the three interior angles of the unitarity triangle [1]. If the measured values of these angles are inconsistent with the predictions of the SM, this will indicate the presence of new physics (NP).

The most promising modes for measuring the CP phases are those that are dominated by a single decay amplitude. In this case, the weak-phase information can be extracted cleanly, i.e. with no hadronic uncertainties. An example of such a decay is the so-called “gold-plated” mode $B_d^0(t) \rightarrow J/\psi K_s$, which is used to probe β ⁴. Note that the decay $B_d^0(t) \rightarrow J/\psi K^*$ is equally gold-plated. The only difference, in comparison to $B_d^0(t) \rightarrow J/\psi K_s$, is that here the final state consists of two vector particles. In this case, one has to do an angular analysis to separate out the CP-even and CP-odd components [1]. Each component can then be treated separately, and β can be obtained cleanly.

Suppose now that there is new physics. How does this affect the above analysis? If the NP affects $B_d^0\text{--}\bar{B}_d^0$ mixing only, the above analysis is unchanged, except that the measured value of β is not the true SM value, but rather one that has been shifted by a new-physics phase. On the other hand, if the NP affects the decay amplitude [3], then the extraction of β is no longer clean – it may be contaminated by hadronic uncertainties. It is this situation that interests us in this paper.

New physics can affect the decay amplitude either at loop level (i.e. in the $b \rightarrow s$ penguin amplitude) or at tree level. Examples of such new-physics models include non-minimal supersymmetric models and models with Z -mediated flavor-changing neutral currents [4]. In all cases, if the new contributions have a different weak phase than that of the SM amplitude, then the measured value of β , β^{meas} , no longer corresponds to the phase of $B_d^0\text{--}\bar{B}_d^0$ mixing, β^{mix} . (Note that β^{mix} could include NP contributions to the mixing.)

If new-physics contributions to the decay amplitude are present, how will we see them? One way is to note that, to a good approximation, β can also be obtained in the SM from $B_d^0(t) \rightarrow \phi K_s$ and $B_d^0(t) \rightarrow \eta' K_s$ [5]. Indeed, at present there appears to be a discrepancy between the value of β extracted from $B_d^0(t) \rightarrow J/\psi K_s$ and that obtained from $B_d^0(t) \rightarrow \phi K_s$ [6]. Should this difference remain as more data is taken, it would provide indirect evidence for a NP amplitude in $B \rightarrow \phi K$.

Still, even in this case, it would be preferable to have *direct* evidence for this new amplitude. Furthermore, if present, we would like to obtain information about it (magnitude, weak and strong phases). It is therefore important to have as many

⁴In fact, there are two weak amplitudes that can contribute to $B_d^0(t) \rightarrow J/\psi K_s$: the tree amplitude and the $b \rightarrow s$ penguin amplitude. However, the weak phases of these two amplitudes are equal (they vanish in the Wolfenstein parametrization of the CKM matrix [2]), so that there is effectively only a single weak amplitude contributing to $B_d^0 \rightarrow J/\psi K_s$. Thus, the extraction of the CP phase β from this decay mode is extremely clean.

independent tests as possible for NP. One possibility is to search for direct CP violation. However, direct CP asymmetries vanish if the strong phase difference between the SM and NP amplitudes is zero, which may well be the case in B decays. (It has been argued that all strong phases in B decays should be quite small, due to the fact that the b -quark is rather heavy.)

In this Letter we show that an angular analysis of B -meson decays to two vector mesons, such as $B_d^0(t) \rightarrow J/\psi K^*$ or ϕK^* , can provide many signals for NP, including several that are nonzero even if the strong phase differences vanish. Furthermore, if *any* NP signal is found, this analysis allows one to place a lower bound on the size of the NP amplitude, and on the difference $|\beta^{meas} - \beta^{mix}|$. As we will see, the analysis can even be used within the SM to analyze decays such as $B_d^0(t) \rightarrow D^{*-} D^{*-}$, from which CP phases cannot be extracted cleanly due to penguin “pollution.”

Consider a $B \rightarrow V_1 V_2$ decay for which a single weak decay amplitude contributes within the SM. Suppose that there is a new-physics amplitude, with a different weak phase, that contributes to the decay. The decay amplitude for each of the three possible helicity states may be generally written as

$$\begin{aligned} A_\lambda &\equiv \text{Amp}(B \rightarrow V_1 V_2)_\lambda = a_\lambda e^{i\delta_\lambda^a} + b_\lambda e^{i\phi} e^{i\delta_\lambda^b}, \\ \bar{A}_\lambda &\equiv \text{Amp}(\bar{B} \rightarrow \bar{V}_1 \bar{V}_2)_\lambda = a_\lambda e^{i\delta_\lambda^a} + b_\lambda e^{-i\phi} e^{i\delta_\lambda^b}, \end{aligned} \quad (1)$$

where a_λ and b_λ represent the SM and NP amplitudes, respectively, ϕ is the new-physics weak phase, the $\delta_\lambda^{a,b}$ are the strong phases, and the helicity index λ takes the values $\{0, \parallel, \perp\}$. Using CPT invariance, the full decay amplitudes can be written as

$$\begin{aligned} \mathcal{A} &= \text{Amp}(B \rightarrow V_1 V_2) = A_0 g_0 + A_\parallel g_\parallel + i A_\perp g_\perp, \\ \bar{\mathcal{A}} &= \text{Amp}(\bar{B} \rightarrow \bar{V}_1 \bar{V}_2) = \bar{A}_0 g_0 + \bar{A}_\parallel g_\parallel - i \bar{A}_\perp g_\perp, \end{aligned} \quad (2)$$

where the g_λ are the coefficients of the helicity amplitudes written in the linear polarization basis. The g_λ depend only on the angles describing the kinematics [7]. For $B = B_d^0$, the above equations enable us to write the time-dependent decay rates as

$$\Gamma(\overline{B_d^0}(t) \rightarrow V_1 V_2) = e^{-\Gamma t} \sum_{\lambda \leq \sigma} \left(\Lambda_{\lambda\sigma} \pm \Sigma_{\lambda\sigma} \cos(\Delta M t) \mp \rho_{\lambda\sigma} \sin(\Delta M t) \right) g_\lambda g_\sigma. \quad (3)$$

Thus, by performing a time-dependent angular analysis of the decay $B_d^0(t) \rightarrow V_1 V_2$, one can measure 18 observables. These are:

$$\begin{aligned} \Lambda_{\lambda\lambda} &= \frac{1}{2}(|A_\lambda|^2 + |\bar{A}_\lambda|^2), \quad \Sigma_{\lambda\lambda} = \frac{1}{2}(|A_\lambda|^2 - |\bar{A}_\lambda|^2), \\ \Lambda_{\perp i} &= -\text{Im}(A_\perp A_i^* - \bar{A}_\perp \bar{A}_i^*), \quad \Lambda_{\parallel 0} = \text{Re}(A_\parallel A_0^* + \bar{A}_\parallel \bar{A}_0^*), \\ \Sigma_{\perp i} &= -\text{Im}(A_\perp A_i^* + \bar{A}_\perp \bar{A}_i^*), \quad \Sigma_{\parallel 0} = \text{Re}(A_\parallel A_0^* - \bar{A}_\parallel \bar{A}_0^*), \end{aligned}$$

$$\begin{aligned}
\rho_{\perp i} &= \text{Re}\left(\frac{q}{p}[A_{\perp}^* \bar{A}_i + A_i^* \bar{A}_{\perp}]\right), \quad \rho_{\perp\perp} = \text{Im}\left(\frac{q}{p} A_{\perp}^* \bar{A}_{\perp}\right), \\
\rho_{\parallel 0} &= -\text{Im}\left(\frac{q}{p}[A_{\parallel}^* \bar{A}_0 + A_0^* \bar{A}_{\parallel}]\right), \quad \rho_{ii} = -\text{Im}\left(\frac{q}{p} A_i^* \bar{A}_i\right),
\end{aligned} \tag{4}$$

where $i = \{0, \parallel\}$. In the above, $q/p = \exp(-2i\beta^{mix})$, where β^{mix} is the weak phase describing $B_d^0 - \bar{B}_d^0$ mixing. Note that β^{mix} may include NP effects (in the SM, $\beta^{mix} = \beta$). Note also that the signs of the various ρ terms depend on the CP-parity of the various helicity states. We have chosen the sign of ρ_{00} and $\rho_{\parallel\parallel}$ to be -1 , which corresponds to the final state $J/\psi K^*$.

The 18 observables given above can be written in terms of 13 theoretical parameters: three a_{λ} 's, three b_{λ} 's, β^{mix} , ϕ , and five strong phase differences defined by $\delta_{\lambda} \equiv \delta_{\lambda}^b - \delta_{\lambda}^a$, $\Delta_i \equiv \delta_{\perp}^a - \delta_i^a$. The explicit expressions for the observables are as follows:

$$\begin{aligned}
\Lambda_{\lambda\lambda} &= a_{\lambda}^2 + b_{\lambda}^2 + 2a_{\lambda}b_{\lambda} \cos \delta_{\lambda} \cos \phi, \\
\Sigma_{\lambda\lambda} &= -2a_{\lambda}b_{\lambda} \sin \delta_{\lambda} \sin \phi, \\
\Lambda_{\perp i} &= 2[a_{\perp}b_i \cos(\Delta_i - \delta_i) - a_i b_{\perp} \cos(\Delta_i + \delta_{\perp})] \sin \phi, \\
\Lambda_{\parallel 0} &= 2[a_{\parallel}a_0 \cos(\Delta_0 - \Delta_{\parallel}) + a_{\parallel}b_0 \cos(\Delta_0 - \Delta_{\parallel} - \delta_0) \cos \phi \\
&\quad + a_0 b_{\parallel} \cos(\Delta_0 - \Delta_{\parallel} + \delta_{\parallel}) \cos \phi + b_{\parallel}b_0 \cos(\Delta_0 - \Delta_{\parallel} + \delta_{\parallel} - \delta_0)] \sin \phi, \\
\Sigma_{\perp i} &= -2[a_{\perp}a_i \sin \Delta_i + a_{\perp}b_i \sin(\Delta_i - \delta_i) \cos \phi + a_i b_{\perp} \sin(\Delta_i + \delta_{\perp}) \cos \phi \\
&\quad + b_{\perp}b_i \sin(\Delta_i + \delta_{\perp} - \delta_i)] \sin \phi, \\
\Sigma_{\parallel 0} &= 2[a_{\parallel}b_0 \sin(\Delta_0 - \Delta_{\parallel} - \delta_0) - a_0 b_{\parallel} \sin(\Delta_0 - \Delta_{\parallel} + \delta_{\parallel})] \sin \phi, \\
\rho_{ii} &= a_i^2 \sin 2\beta^{mix} + 2a_i b_i \cos \delta_i \sin(2\beta^{mix} + \phi) + b_i^2 \sin(2\beta^{mix} + 2\phi), \\
\rho_{\perp\perp} &= -a_{\perp}^2 \sin 2\beta^{mix} - 2a_{\perp}b_{\perp} \cos \delta_{\perp} \sin(2\beta^{mix} + \phi) - b_{\perp}^2 \sin(2\beta^{mix} + 2\phi), \\
\rho_{\perp i} &= 2[a_i a_{\perp} \cos \Delta_i \cos 2\beta^{mix} + a_{\perp}b_i \cos(\Delta_i - \delta_i) \cos(2\beta^{mix} + \phi) \\
&\quad + a_i b_{\perp} \cos(\Delta_i + \delta_{\perp}) \cos(2\beta^{mix} + \phi) \\
&\quad + b_i b_{\perp} \cos(\Delta_i + \delta_{\perp} - \delta_i) \cos(2\beta^{mix} + 2\phi)] \sin \phi, \\
\rho_{\parallel 0} &= 2[a_0 a_{\parallel} \cos(\Delta_0 - \Delta_{\parallel}) \sin 2\beta^{mix} + a_{\parallel}b_0 \cos(\Delta_0 - \Delta_{\parallel} - \delta_0) \sin(2\beta^{mix} + \phi) \\
&\quad + a_0 b_{\parallel} \cos(\Delta_0 - \Delta_{\parallel} + \delta_{\parallel}) \sin(2\beta^{mix} + \phi) \\
&\quad + b_0 b_{\parallel} \cos(\Delta_0 - \Delta_{\parallel} + \delta_{\parallel} - \delta_0) \sin(2\beta^{mix} + 2\phi)] \sin \phi.
\end{aligned} \tag{5}$$

It is straightforward to show that one cannot extract β^{mix} . There are a total of six amplitudes describing $B \rightarrow V_1 V_2$ and $\bar{B} \rightarrow \bar{V}_1 \bar{V}_2$ decays [Eq. (1)]. Thus, at best one can measure the magnitudes and relative phases of these six amplitudes, giving 11 measurements. Since the number of measurements (11) is fewer than the number of theoretical parameters (13), one cannot obtain any of the theoretical unknowns purely in terms of observables. In particular, it is impossible to extract β^{mix} cleanly.

In the absence of NP, $b_\lambda = 0$. The number of parameters is then reduced from 13 to 6: three a_λ 's, two strong phase differences (Δ_i), and β^{mix} . All of these can be determined cleanly in terms of observables. Because we have 18 observables, but only 6 theoretical parameters, there are 12 relations that must exist among the observables in the absence of NP. (Of course, only five of these are independent.) The 12 relations are:

$$\begin{aligned}
\Sigma_{\lambda\lambda} &= \Lambda_{\perp i} = \Sigma_{\parallel 0} = 0 \\
\frac{\rho_{ii}}{\Lambda_{ii}} &= -\frac{\rho_{\perp\perp}}{\Lambda_{\perp\perp}} = \frac{\rho_{\parallel 0}}{\Lambda_{\parallel 0}} \\
\Lambda_{\parallel 0} &= \frac{1}{2\Lambda_{\perp\perp}} \left[\frac{\Lambda_{\lambda\lambda}^2 \rho_{\perp 0} \rho_{\perp\parallel} + \Sigma_{\perp 0} \Sigma_{\perp\parallel} (\Lambda_{\lambda\lambda}^2 - \rho_{\lambda\lambda}^2)}{\Lambda_{\lambda\lambda}^2 - \rho_{\lambda\lambda}^2} \right] \\
\frac{\rho_{\perp i}^2}{4\Lambda_{\perp\perp}\Lambda_{ii} - \Sigma_{\perp i}^2} &= \frac{\Lambda_{\perp\perp}^2 - \rho_{\perp\perp}^2}{\Lambda_{\perp\perp}^2}.
\end{aligned} \tag{6}$$

The key point is the following: *the violation of any of the above relations will be a smoking-gun signal of NP*. We therefore see that the angular analysis of $B \rightarrow V_1 V_2$ decays provides numerous tests for the presence of new physics.

The observable $\Lambda_{\perp i}$ deserves special attention [8]. From Eq. (5), one sees that even if the strong phase differences vanish, $\Lambda_{\perp i}$ is nonzero in the presence of new physics ($\phi \neq 0$), in stark contrast to the direct CP asymmetries (proportional to $\Sigma_{\lambda\lambda}$). This is due to the fact that the $\perp$ helicity is CP-odd, while the 0 and $\parallel$ helicities are CP-even. Thus, $\perp$ -0 and $\perp$ - $\parallel$ interferences include an additional factor of ' i ' in the full decay amplitudes [Eq. (2)], which leads to the cosine dependence on the strong phases.

Now, although the reconstruction of the full $B_d^0(t)$ and $\bar{B}_d^0(t)$ decay rates in Eq. (3) requires both tagging and time-dependent measurements, the $\Lambda_{\lambda\sigma}$ terms remain even if the two rates for $B_d^0(t)$ and $\bar{B}_d^0(t)$ decays are added together. Note also that these terms are time-independent. Therefore, *no tagging or time-dependent measurements are needed to extract $\Lambda_{\perp i}$!* It is only necessary to perform an angular analysis of the final state $V_1 V_2$. Thus, this measurement can even be made at a symmetric B -factory.

The decays of charged B mesons to vector-vector final states are even simpler to analyze since no mixing is involved. One can in principle combine charged and neutral B decays to increase the sensitivity to new physics. For example, for $B \rightarrow J/\psi K^*$ decays, one simply performs an angular analysis on all decays in which a J/ψ is produced accompanied by a charged or neutral K^* . A nonzero value of $\Lambda_{\perp i}$ would be a clear signal for new physics [9].

The decays of both charged and neutral B mesons to $D_s^* D^*$ can be analyzed similarly. Because these modes are dominated by a single decay amplitude in the SM, no direct CP violation is expected. And since this is not a final state to which both B_d^0 and $\bar{B}_d^0$ can decay, no indirect CP violation is present either. An angular

analysis of these decays would therefore be interesting – if any CP-violating signal were found, such as a nonzero value of $\Lambda_{\perp i}$, this would again indicate the presence of new physics.

It must be noted that, despite the large number of new-physics signals, it is still possible for the NP to remain hidden. This happens if a singular situation is realized. If the three strong phase differences δ_λ vanish, and the ratio $r_\lambda \equiv b_\lambda/a_\lambda$ is the same for all helicities, i.e. $r_0 = r_\parallel = r_\perp$, then it is easy to show that the relations in Eq. (6) are all satisfied. Thus, if the NP happens to respect these very special conditions, the angular analysis of $B \rightarrow V_1 V_2$ would show no signal for NP even if it is present, and the measured value of β would not correspond to the actual phase of $B_d^0 - \bar{B}_d^0$ mixing.

Now, suppose that some signal of new physics is found, indicating that β^{meas} is not equal to β^{mix} . As we have argued earlier, in the presence of new physics one cannot extract the true value of β^{mix} . However, as we will describe below, the angular analysis does allow one to constrain the value of the difference $|\beta^{meas} - \beta^{mix}|$, as well as the size of the NP amplitudes b_λ^2 . Naively, one would not think it possible to obtain any constraints on the NP parameters. After all, we have 11 measurements, but 13 theoretical unknown parameters. However, because the equations are nonlinear, such constraints are possible. Below, we list some of these constraints; their full derivation will be presented elsewhere [10].

In the constraints, we will make use of the following quantities. For the vector-vector final state, the analogue of the usual direct CP asymmetry a_{dir}^{CP} is $a_\lambda^{dir} \equiv \Sigma_{\lambda\lambda}/\Lambda_{\lambda\lambda}$, which is helicity-dependent. For convenience, we define the related quantity $y_\lambda = \sqrt{1 - \Sigma_{\lambda\lambda}^2/\Lambda_{\lambda\lambda}^2}$. The measured value of $\sin 2\beta$ can also depend on the helicity of the final state: $\rho_{\lambda\lambda}$ can be recast in terms of a measured weak phase $2\beta_\lambda^{meas}$, defined as

$$\sin 2\beta_\lambda^{meas} = \frac{\pm \rho_{\lambda\lambda}}{\sqrt{\Lambda_{\lambda\lambda}^2 - \Sigma_{\lambda\lambda}^2}}, \quad (7)$$

where the $+$ ($-$) sign corresponds to $\lambda = 0, \parallel$ ($\perp$). In terms of these quantities, the size of NP amplitudes b_λ^2 may be expressed as

$$2b_\lambda^2 \sin^2 \phi = \Lambda_{\lambda\lambda} (1 - y_\lambda \cos(2\beta_\lambda^{meas} - 2\beta)). \quad (8)$$

The form of the constraints depends on which new-physics signals are observed; we give a partial list below. For example, suppose that direct CP violation is observed in a particular helicity state. In this case a lower bound on the corresponding NP amplitude can be obtained by minimizing b_λ^2 [Eq. (8)] with respect to β and ϕ :

$$b_\lambda^2 \geq \frac{1}{2} \Lambda_{\lambda\lambda} [1 - y_\lambda]. \quad (9)$$

On the other hand, suppose that the new-physics signal is $\beta_i^{meas} \neq \beta_j^{meas}$. Defining $2\omega \equiv 2\beta_j^{meas} - 2\beta_i^{meas}$ and $\eta_\lambda \equiv 2(\beta_\lambda^{meas} - \beta^{mix})$, the minimization of $(b_i^2 \mp b_j^2)$ with

respect to η_i and ϕ yields

$$(b_i^2 \mp b_j^2) \geq \frac{\Lambda_{ii} \mp \Lambda_{jj}}{2} - \frac{|y_i \Lambda_{ii} \mp y_j \Lambda_{jj} e^{2i\omega}|}{2}, \quad (10)$$

where $\Lambda_{ii} > \Lambda_{jj}$ is assumed. If there is no direct CP violation ($\Sigma_{\lambda\lambda} = 0$), but $\Lambda_{\perp i}$ is nonzero, one has

$$2(b_i^2 \mp b_{\perp}^2) \geq \Lambda_{ii} \mp \Lambda_{\perp\perp} - \sqrt{(\Lambda_{ii} \mp \Lambda_{\perp\perp})^2 \pm \Lambda_{\perp i}^2}, \quad (11)$$

where $\beta_{\perp}^{meas}$ was eliminated using the expression for $\Lambda_{\perp i}$.

One can also obtain bounds on $|\beta_{\lambda}^{meas} - \beta^{mix}|$, though this requires the nonzero measurement of observables involving the interference of different helicities. For example, if $\Lambda_{\perp i}$ is nonzero and $\Sigma_{\lambda\lambda} = 0$, we find

$$\begin{aligned} \Lambda_{ii} \cos \eta_i + \Lambda_{\perp\perp} \cos(\eta_{\perp} - 2\eta_i) &\leq \sqrt{(\Lambda_{ii} + \Lambda_{\perp\perp})^2 - \Lambda_{\perp i}^2}, \\ \Lambda_{ii} \cos \eta_i - \Lambda_{\perp\perp} \cos \eta_{\perp} &\leq \sqrt{(\Lambda_{ii} - \Lambda_{\perp\perp})^2 + \Lambda_{\perp i}^2}. \end{aligned} \quad (12)$$

If $\Lambda_{\perp i} \neq 0$, one cannot have $\eta_i = \eta_{\perp} = 0$. These constraints therefore place a lower bound on $|\beta_i^{meas} - \beta^{mix}|$ and/or $|\beta_{\perp}^{meas} - \beta^{mix}|$.

A-priori, one does not know which of the above constraints is strongest – this depends on the actual values of the observables. Of course, in practice, one will simply perform a fit to obtain the best lower bounds on these NP parameters [10]. However, it is interesting to see that constraints can be obtained analytically.

As a specific application, we have noted the apparent discrepancy in the value of $\sin 2\beta$ as obtained from measurements of $B_d^0(t) \rightarrow J/\psi K_S$ and $B_d^0(t) \rightarrow \phi K_S$ [6]. In this case, the angular analyses of $B_d^0(t) \rightarrow J/\psi K^*$ and $B_d^0(t) \rightarrow \phi K^*$ would allow one to determine if new physics is indeed present. If NP is confirmed, this analysis would allow one to put constraints on the NP parameters.

Note that this analysis can also be applied within the SM to decays such as $B_d^0(t) \rightarrow D^{*+} D^{*-}$. These decays have both a tree and a penguin contribution, so that β^{mix} cannot be extracted cleanly. Assuming no new physics, the above analysis allows one to obtain lower bounds on the ratio of penguin to tree amplitudes, as well as on $|\beta_{\lambda}^{meas} - \beta^{mix}|$. This can serve as a crosscheck on other measurements of β^{mix} , as well as on model calculations of the hadronic amplitudes.

Finally, it is worthwhile to examine the feasibility of this method. The present data at B -factories can already be used to perform time-independent angular analyses of $B \rightarrow V_1 V_2$ decays. In fact, BaBar has measured a nonzero value of $\Lambda_{\perp i}$ (a CP-violating triple-product correlation) in $B \rightarrow \phi K^*$ at 1.7σ [11]. This is a potential hint of new physics. On the other hand, time-dependent angular analyses will take considerably more time to carry out. Thus, it may be several more years before we have new-physics signals which rely on time-dependent measurements.

To sum up: in the standard model (SM), the cleanest extraction of the CP angles comes from neutral B decays that are dominated by a single decay amplitude. If

there happens to be a new-physics (NP) contribution to the decay amplitude, with a different weak phase, this could seriously affect the cleanliness of the measurement. There is already a hint of such NP, as indicated by the discrepancy between the value of β extracted from $B_d^0(t) \rightarrow J/\psi K_S$ and that obtained from $B_d^0(t) \rightarrow \phi K_S$. However, it is important to confirm this through independent direct tests, and to attempt to obtain information about the NP amplitude, if possible.

In this paper, we have shown that this type of new physics can be probed by performing an angular analysis of the related $B \rightarrow V_1 V_2$ decay modes. There are numerous relations that are violated in the presence of NP, and several of these remain nonzero even if the strong phase difference between the SM and NP amplitudes vanishes. The most incisive test is a measurement of $\Lambda_{\perp i} \neq 0$. To obtain this observable, neither tagging nor time-dependent measurements is necessary – one can combine all neutral and charged B decays.

Furthermore, should a signal for new physics be found, one can place a lower bound on the difference $|\beta^{meas} - \beta^{mix}|$, as well as on the size of the NP amplitudes. By applying this analysis to the decays $B_d^0(t) \rightarrow J/\psi K^*$ and $B_d^0(t) \rightarrow \phi K^*$, one can confirm the presence of the new physics that is hinted at in the measurements of $B_d^0(t) \rightarrow J/\psi K_S$ and $B_d^0(t) \rightarrow \phi K_S$ [6]. It can even be applied within the SM to decays such as $B_d^0(t) \rightarrow D^{*+} D^{*-}$, which receive both tree and penguin contributions.

Acknowledgements: N.S. and R.S. thank D.L. for the hospitality of the Université de Montréal, where part of this work was done. The work of D.L. was financially supported by NSERC of Canada. The work of N.S. was supported by the Department of Science and Technology, India.

References

- [1] For a review, see, for example, *The BaBar Physics Book*, eds. P.F. Harrison and H.R. Quinn, SLAC Report 504, 1998.
- [2] L. Wolfenstein, *Phys. Rev. Lett.* **51**, 1945, 83.
- [3] Y. Grossman and M.P. Worah, *Phys. Lett.* **395B**, 241, 97.
- [4] See, for example, M. Gronau and D. London, *Phys. Rev.* **D55**, 2845, 97; Y. Grossman and M.P. Worah, Ref. [3].
- [5] D. London and A. Soni, *Phys. Lett.* **407B**, 61, 97.
- [6] For a review, see T. Browder, talk given at Lepton-Photon 2003, Fermilab, Batavia, Illinois, 11-16 August 2003, <http://conferences.fnal.gov/lp2003>.
- [7] N. Sinha and R. Sinha, *Phys. Rev. Lett.* **80**, 3706, 98; A.S. Dighe, I. Dunietz and R. Fleischer, *Eur. Phys. J.* **C6**, 647, 99.

- [8] The observables $\Lambda_{\perp i}$, which are triple-product correlations, have also been studied in $B \rightarrow V_1 V_2$ decays in A. Datta and D. London, hep-ph/0303159.
- [9] This point has also been made in X.-G. He and W.-S. Hou, *Phys. Rev.* **D58**: 117502, 1998; D. London, N. Sinha and R. Sinha, hep-ph/0207007.
- [10] D. London, N. Sinha and R. Sinha, arXiv:hep-ph/0402214.
- [11] Talk by Jim Smith at Moriond, 2004:
<http://moriond.in2p3.fr/QCD/2004/TuesdayMorning/Smith.pdf>.